

Package ‘cramer’

May 8, 2026

Version 0.9-4

Date 2024-02-04

Title Multivariate Nonparametric Cramer-Test for the
Two-Sample-Problem

Author Carsten Franz <carsten.franz@gmail.com>

Maintainer Carsten Franz <carsten.franz@gmail.com>

Depends R (>= 0.65), boot

Description Provides R routine for the so called two-sample
Cramer-Test. This nonparametric two-sample-test on equality
of the underlying distributions can be applied to
multivariate data as well as univariate data. It offers two
possibilities to approximate the critical value both of which
are included in this package.

License GPL (>= 2)

Repository CRAN

Encoding UTF-8

Imports Rcpp (>= 1.0.12)

LinkingTo Rcpp

RoxygenNote 7.3.1

NeedsCompilation yes

Date/Publication 2024-02-05 21:30:02 UTC

Contents

cramer-package	2
cramer.test	2

Index	6
--------------	----------

 cramer-package

Cramer-Test for uni- and multivariate two-sample-problem

Description

Provides R routine for the so called two-sample Cramer-Test. This nonparametric two-sample-test on equality of the underlying distributions can be applied to multivariate data as well as univariate data. It offers two possibilities to approximate the critical value both of which are included in this package.

Author(s)

Carsten Franz <carsten.franz@gmail.com>

 cramer.test

Perform Cramer-Test for uni- and multivariate two-sample-problem

Description

Perform Cramér-test for two-sample-problem.

Both univariate and multivariate data is possible. For calculation of the critical value Monte-Carlo-bootstrap-methods and eigenvalue-methods are available. For the bootstrap access ordinary and permutation methods can be chosen as well as the number of bootstrap-replicates taken.

Usage

```
cramer.test(
  x,
  y,
  conf.level = 0.95,
  replicates = 1000,
  sim = "ordinary",
  just.statistic = FALSE,
  kernel = "phiCramer",
  maxM = 2^14,
  K = 160
)
```

Arguments

x	First set of observations. Either in vector form (univariate) or in a matrix with one observation per row (multivariate).
y	Second set of observations. Same dimension as x.
conf.level	Confidence level of test. The default is conf.level=0.95.

replicates	Number of bootstrap-replicates taken to obtain critical value. The default is replicates=1000. When using the eigenvalue method, this variable is unused.
sim	Type of Monte-Carlo-bootstrap method or eigenvalue method. Possible values are "ordinary" (default) for normal Monte-Carlo-bootstrap, "permutation" for a permutation Monte-Carlo-bootstrap or "eigenvalue" for bootstrapping the limit distribution, evaluating the (approximate) eigenvalues being the weights of the limiting chisquared-distribution and using the critical value of this approximation (calculated via fast fourier transform). This method is especially good if the dataset is too large to perform Monte-Carlo-bootstrapping (although it must not be too large so the matrix eigenvalue problem can still be solved).
just.statistic	Boolean variable. If TRUE just the value of the Cramér-statistic is calculated and no bootstrap-replicates are produced.
kernel	Character-string giving the name of the kernel function. The default is "phiCramer" which is the Cramér-test included in earlier versions of this package and which is used in the paper of Baringhaus and the author mentioned below. It is possible to use user-defined kernel functions here. The functions needs to be able to deal with matrix arguments. Kernel functions need to be defined on the positive real line with value 0 at 0 and have a non-constant completely monotone first derivative. An example is show in the Examples section below. Build-in functions are "phiCramer", "phiBahr", "phiLog", "phiFracA" and "phiFracB".
maxM	Gives the maximum number of points used for the fast fourier transform. When using Monte-Carlo-bootstrap methods, this variable is unused.
K	Gives the upper value up to which the integral for the calculation of the distribution function out of the characteristic function (Gurlands formula) is evaluated. The default ist 160. Careful: When increasing K it is necessary to increase maxM as well since the resolution of the points where the distribution function is calculated is

$$\frac{2\pi}{K}.$$

Thus, if just K is increased the maximum value, where the distribution function is calculated is lower. When using Monte-Carlo-bootstrap methods, this variable is unused.

Details

The Cramér-statistic is given by

$$T_{m,n} = \frac{mn}{m+n} \left(\frac{2}{mn} \sum_{i,j}^{m,n} \phi(\|\vec{X}_i - \vec{Y}_j\|^2) - \frac{1}{m^2} \sum_{i,j=1}^m \phi(\|\vec{X}_i - \vec{X}_j\|^2) - \frac{1}{n^2} \sum_{i,j=1}^n \phi(\|\vec{Y}_i - \vec{Y}_j\|^2) \right),$$

The function ϕ is the kernel function mentioned in the Parameters section. The proof that the Monte-Carlo-Bootstrap and eigenvalue methods work is given in the reference given below. Other build-in kernel functions are

$$\phi_{Cramer}(z) = \sqrt{z}/2$$

(recommended for location alternatives),

$$\phi_{Bahrr}(z) = 1 - \exp(-z/2)$$

(recommended for dispersion as well as location alternatives),

$$\phi_{log}(z) = \log(1 + z)$$

(preferrably for location alternatives),

$$\phi_{FracA}(z) = 1 - \frac{1}{1 + z}$$

(preferrably for dispersion alternatives) and

$$\phi_{FracB}(z) = 1 - \frac{1}{(1 + z)^2}.$$

(also for dispersion alternatives). Test performance was investigated in the below referenced 2010 publication. The idea of using this statistic is due to L. Baringhaus, University of Hanover.

Value

The returned value is an object of class "cramertest", containing the following components:

method	Describing the test in words.
d	Dimension of the observations.
m	Number of x observations.
n	Number of y observations.
statistic	Value of the Cramér-statistic for the given observations.
conf.level	Confidence level for the test.
crit.value	Critical value calculated by bootstrap method, eigenvalue method, respectively. When using the eigenvalue method, the distribution under the hypothesis will be interpolated linearly.
p.value	Estimated p-value of the test.
result	Contains 1 if the hypothesis of equal distributions should not be accepted and 0 otherwise.
sim	Method used for obtaining the critical value.
replicates	Number of bootstrap-replicates taken.
ev	Contains eigenvalues and eigenfunctions when using the eigenvalue-method to obtain the critical value
hypdist	Contains the via fft reconstructed distribution function under the hypothesis. \$x contains the x-values and \$Fx the values of the distribution function at the positions.

References

The test and its properties is described in:

- Baringhaus, L. and Franz, C. (2004) *On a new multivariate two-sample test*, Journal of Multivariate Analysis, 88, p. 190-206
- Baringhaus, L. and Franz, C. (2010) *Rigid motion invariant two-sample tests*, Statistica Sinica 20, 1333-1361

The test of Bahr is also discussed in:

- Bahr, R. (1996) *Ein neuer Test fuer das mehrdimensionale Zwei-Stichproben-Problem bei allgemeiner Alternative*, German, Ph.D. thesis, University of Hanover

Examples

```
# comparison of two univariate normal distributions
x<-rnorm(20,mean=0,sd=1)
y<-rnorm(50,mean=0.5,sd=1)
cramer.test(x,y)

# comparison of two multivariate normal distributions with permutation test:
# library "MASS" for multivariate routines (included in package "VR")

# library(MASS)
# x<-mvrnorm(n=20,mu=c(0,0),Sigma=diag(c(1,1)))
# y<-mvrnorm(n=50,mu=c(0.3,0),Sigma=diag(c(1,1)))
# cramer.test(x,y,sim="permutation")

# comparison of two univariate normal distributions with Bahrs Kernel
phiBahr<-function(x) return(1-exp(-x/2))
x<-rnorm(20,mean=0,sd=1)
y<-rnorm(50,mean=0,sd=2)
cramer.test(x,y,sim="eigenvalue",kernel="phiBahr")
```

Index

cramer (cramer-package), [2](#)
cramer-package, [2](#)
cramer.test, [2](#)