

Package ‘drcSeedGerm’

May 8, 2026

Date 2023-04-20

Type Package

Title Utilities for Data Analyses in Seed Germination/Emergence Assays

Version 1.0.1

Depends R (>= 3.5.0), drc, drcte

Imports plyr, dplyr, mvtnorm, survival

Author Andrea Onofri <andrea.onofri@unipg.it>

Maintainer Andrea Onofri <andrea.onofri@unipg.it>

Description Utility functions to be used to analyse datasets obtained from seed germination/emergence assays. Fits several types of seed germination/emergence models, including those reported in Onofri et al. (2018) ``Hydrothermal-time-to-event models for seed germination'', European Journal of Agronomy, 101, 129-139 <doi:10.1016/j.eja.2018.08.011>. Contains several datasets for practicing.

URL <https://www.statforbiology.com>

License GPL (>= 2)

Encoding UTF-8

LazyData true

RoxygenNote 7.2.0

NeedsCompilation no

Repository CRAN

Date/Publication 2023-05-16 08:30:05 UTC

Contents

alfalfaSG	2
barley	4
excalibur	5
festuca	6
GR_Psi_models	7
GR_TPsi_models	8

GR_T_models	10
hordeum	11
HTE1	13
HTE2	14
HTE3	16
HTex	18
HTG	19
HTL	21
HTLL	22
HTnorm	24
HTTEM	25
HTTLL	27
HTTnorm.M	29
HTW1	31
HTW2	32
jackGroupSE	34
makeDrm	35
makeDrm2	35
phalaris	36
Pmax_models	38
rape	39
rape2G	40
Thermal-Time_models	41
Index	44

 alfalfaSG

Field-book for a germination assay with alfalfa

Description

The germination of alfalfa was assayed at 7 temperature levels, on three replicated Petri dishes per temperature and 100 seeds per Petri dish. Inspections were made in several times after the beginning of the assay.

Usage

```
data("alfalfaSG")
```

Format

A data frame with 21 observations on the following variables.

Dish a numeric vector with the coding for Petri dishes

Temp a numeric vector with the temperature level

nViable a numeric vector: number of viable seeds per dish

'1' a numeric vector: count at day 1

'2' a numeric vector: count at day 2
'3' a numeric vector: count at day 3
'4' a numeric vector: count at day 4
'5' a numeric vector: count at day 5
'6' a numeric vector: count at day 6
'7' a numeric vector: count at day 7
'8' a numeric vector: count at day 8
'9' a numeric vector: count at day 9
'10' a numeric vector: count at day 10
'11' a numeric vector: count at day 11
'12' a numeric vector: count at day 12
'13' a numeric vector: count at day 13
'14' a numeric vector: count at day 14
'15' a numeric vector: count at day 15
'16' a numeric vector: count at day 16
'17' a numeric vector: count at day 17
'18' a numeric vector: count at day 18
'19' a numeric vector: count at day 19
'20' a numeric vector: count at day 20
'21' a numeric vector: count at day 21
'22' a numeric vector: count at day 22
'23' a numeric vector: count at day 23
'24' a numeric vector: count at day 24
'27' a numeric vector: count at day 27
'28' a numeric vector: count at day 28
'29' a numeric vector: count at day 29
'31' a numeric vector: count at day 31
'34' a numeric vector: count at day 34

Details

Every line of data represents a Petri dish. There were 100 seeds per Petri dish. The columns represent the characteristics of each Petri dish. The columns from 4 to 32 represent the number of germinated seeds counted at each assessment time.

Author(s)

Andrea Onofri

Source

no reference yet

References

Onofri, A., Benincasa, P., Mesgaran, M.B., Ritz, C., 2018. Hydrothermal-time-to-event models for seed germination. *European Journal of Agronomy* 101, 129–139.

Examples

```
data(alfalfaSG)
head(alfalfaSG)
```

barley	<i>A series of germination assays with barley</i>
--------	---

Description

The germination of barley was assayed at 9 temperature levels, on three replicated Petri dishes at each temperature and 50 seeds per Petri dish. Inspections were made in several times after the beginning of the assay.

Usage

```
data("barley")
```

Format

A data frame with 810 observations on the following 7 variables.

Dish a numeric vector with the coding for Petri dishes

Temp a numeric vector with the temperature level

timeBef a numeric vector, with the start time for each inspection interval

timeAf a numeric vector, with the end time for each inspection interval

nSeeds a numeric vector, with the number of germinated seeds at each inspection interval

nCum a numeric vector, with the cumulative number of germinated seeds at each assessment time

propCum a numeric vector, the cumulative proportion of germinated seeds at each assessment time

Details

The variable 'timeAf' contains the value 'Inf' (Infinity), that corresponds to the seeds which did not germinate during the assay, for which we the germination time might be comprised from the last assessment time to infinity.

Author(s)

Andrea Onofri

Source

no reference yet

References

Onofri, A., Benincasa, P., Mesgaran, M.B., Ritz, C., 2018. Hydrothermal-time-to-event models for seed germination. *European Journal of Agronomy* 101, 129–139.

Examples

```
data(barley)
```

excalibur	<i>Relationship between germination rate and water potential in oilseed rape (var. Excalibur)</i>
-----------	---

Description

This files describes the relationship between germination rate and water potential in the substrate for seeds of oilseed rape (var. Excalibur). Three germination percentiles are considered for germination rate, i.e. GR10, GR30 and GR50.

Usage

```
data("excalibur")
```

Format

A data frame with 27 observations on the following 5 variables.

Perc a numeric vector: the germination percentile

Psi a numeric vector: water potential in the substrate (in MPa)

Tg a numeric vector: germination time in days

SE a numeric vector: standard errors for germination times (in days)

GR a numeric vector: germination rates in d^{-1}

Source

Pace, R., Benincasa, P., Ghanem, M.E., Quinet, M., Lutts, S., 2012. GERMINATION OF UNTREATED AND PRIMED SEEDS IN RAPESEED (BRASSICA NAPUS VAR OLEIFERA DEL.) UNDER SALINITY AND LOW MATRIC POTENTIAL. *Experimental Agriculture* 48, 238–251.

References

Pace, R., Benincasa, P., Ghanem, M.E., Quinet, M., Lutts, S., 2012. Germination of untreated and primed seeds in rapeseed (*Brassica napus* var. *oleifera* Del.) under salinity and low matric potential. *Experimental Agriculture* 48, 238–251.

Examples

```

data(excalibur)
head(excalibur)
library(drc)

modGR1 <- drm(GR ~ Psi, fct=GRPsiLin(), data=excalibur, curveid=Perc)
summary(modGR1)
plot(modGR1, log="", legendPos=c(-0.9, 1))

modGR2 <- drm(GR ~ Psi, fct=GRPsiPol2(), data=excalibur, curveid=Perc)
summary(modGR2)
plot(modGR2, log="", legendPos=c(-0.9, 1))

```

festuca

Relationship between germination rate and water potential in Festuca arundinacea L.

Description

This files describes the relationship between germination rate and Festuca arundinacea. Three germination percentiles are considered for germination rate, i.e. GR10, GR30 and GR50.

Usage

```
data("festuca")
```

Format

A data frame with 36 observations on the following 3 variables.

g a numeric vector: the germination percentile

Psi a numeric vector: water potential in the substrate (in MPa)

GR a numeric vector: germination rates in d^{-1}

Source

TEI F, BENINCASA P and CIRICIOFOLO E (2001) Effetto del potenziale idrico e della temperatura sulla germinazione di alcune specie graminacee da tappeto erboso. In: Atti XXXIV Convegno della Società Italiana di Agronomia, Pisa, Italy, 200–201.

References

TEI F, BENINCASA P and CIRICIOFOLO E (2001) Effetto del potenziale idrico e della temperatura sulla germinazione di alcune specie graminacee da tappeto erboso. In: Atti XXXIV Convegno della Società Italiana di Agronomia, Pisa, Italy, 200–201.

Examples

```
data(festuca)
modGR1 <- drm(GR ~ Psi, fct=GRPsiLin(),
              data=festuca, curveid=g)
summary(modGR1)
```

GR_Psi_models

*Hydro-time models based on the germination rate***Description**

These models describe the response of germination rate to water potential in the substrate.

Usage

```
GRPsiLin()
GRPsiPol()
GRPsiPol2()
GRPsi.Lin()
GRPsi.Pol()
GRPsi.Pol2()
GRPsiLin.fun(Psi, Psib, thetaH)
GRPsiPol.fun(Psi, Psib, thetaH)
GRPsiPol2.fun(Psi, Psib, thetaH)
```

Arguments

The functions 'GRPsiLin()', 'GRPsiPol()', 'GRPsiPol2()', GRPsi.Lin(), GRPsi.Pol(), GRPsi.Pol2() have no arguments. The functions 'GRPsiLin.fun()', 'GRPsiPol.fun()' and 'GRPsiPol2.fun()' have the following arguments:

Psi	Water potential in the substrate
Psib	Base water potential within the population
thetaH	Hydro-time parameter

Details

The functions 'GRPsiLin()', 'GRPsiPol()', 'GRPsiPol2()', GRPsi.Lin(), GRPsi.Pol(), GRPsi.Pol2() are meant to be used with the 'drm()' function in the 'drc' package ('GRPsiLin()' and 'GRPsi.Lin()', 'GRPsiPol()' and 'GRPsi.Pol()', 'GRPsiPol2()' and 'GRPsiPol2()') are totally equivalent, apart from the names). The functions 'GRPsiLin.fun()', 'GRPsiPol.fun()' and 'GRPsiPol2.fun()' are used for general purposes (plotting and other uses). Details about these functions and the meaning of parameters are described in Bradford (2002) and in the package documentation (see references below).

Value

The 'GRPsiLin.fun()', 'GRPsiPol.fun()' and 'GRPsiPol2.fun()' functions return the germination rate for any given values of water potential in the substrate. The 'GRPsiLin()', 'GRPsiPol()' and 'GRPsiPol2()' (and 'GRPsi.Lin()', 'GRPsi.Pol()' and 'GRPsi.Pol2()') functions return a list containing the nonlinear function, the self starter function, the parameter names and other items which are internally used by the 'drc()' function.

Author(s)

Andrea Onofri

References

See package documentation at: https://www.statforbiology.com/_seedtutorial/

Examples

```
library(drcte)
# Observed data
Psi <- c(-2, -1.5, -1.2, -1, -0.8, -0.6, -0.4, -0.25,
        -0.12, -0.06, -0.03, 0)
GR <- c(0, 0, 0, 0, 0.0585, 0.094, 0.1231, 0.1351,
        0.1418, 0.1453, 0.1458, 0.1459)
Psi2 <- c(-0.5, -0.6, -0.7, -0.8, -0.9, -1, -1.1, -1.2,
        -1.5)
GR2 <- c(1.4018, 1.0071, 0.5614, 0.3546, 0.2293, 0, 0,
        0, 0)

# Model fitting
modHT1 <- drm(GR ~ Psi, fct = GRPsiLin())
modHT2 <- drm(GR ~ Psi, fct = GRPsiPol())
modHT3 <- drm(GR2 ~ Psi2, fct = GRPsiPol2())

summary(modHT1)
summary(modHT2)
summary(modHT2)
```

Description

These models are used to describe the germination rate of a seed, depending on the environmental temperature and water potential.

Usage

```
GRTPsi.M()
GRTPsi.M.fun(Temp, Psi, k, Tb, ThetaHT, Psib)
GRTPsi.BS()
GRTPsi.BS.fun(Temp, Psi, k, Tb, To, ThetaHT, Psib)
```

Arguments

Temp	Temperature variable
Psi	Water potential variable
Psib	Base water potential variable
k	regression parameter
Tb	base temperature
To	optimal temperature
ThetaHT	Hydro-thermal-time parameter

Details

The 'GRT.M.fun()' is a generic R function, while the GRT.M() function is meant to be used with the 'drm()' function, within the 'drc' package.

Value

The 'GRT.M.fun()' functions returns a vector of responses, for given values of temperature, Tc, Tb and ThetaH. The GRT.M() function returns a list containing the nonlinear function, the self starter function, the parameter names and other items which are internally used by the 'drm()' function.

Author(s)

Andrea Onofri

References

https://www.statforbiology.com/2023/stat_drcte_12-htt2step/

Examples

```
library(drc)
Tval <- c(2, 5, 10, 15, 20, 25, 30, 35, 40)
GR <- c(0, 0, 0.209, 0.435, 0.759, 0.821, 0.417, 0.145, 0)
modM <- drm(GR ~ Tval, fct = GRT.M())
plot(modM, log="", xlim = c(0, 40), ylim=c(0,1.2),
      legendPos = c(5, 1.0), xlab = "Temperature (°C)")
```

Description

These models are used to describe the germination rate of a seed, depending on the environmental temperature.

Usage

```
GRT.GH()
GRT.GH2()
GRT.YL()
GRT.BS()
GRT.BSb()
GRT.Ex()
GRT.Exb()
GRT.M()
GRT.Mb()
GRT.RF()
GRT.RFb()
GRT.GH.fun(Temp, Tb, ThetaT)
GRT.GH2.fun(Temp, Tb, beta)
GRT.YL.fun
GRT.BS.fun(Temp, k, Tb, To, ThetaT)
GRT.BSb.fun(Temp, Tc, Tb, To, ThetaT)
GRT.Ex.fun(Temp, k, Tb, Tc, ThetaT)
GRT.Exb.fun(Temp, k, Tb, Tc, ThetaT)
GRT.M.fun(Temp, k, Tb, ThetaT)
GRT.Mb.fun(Temp, Tb, Tc, ThetaT)
GRT.RF.fun(Temp, k, Tb, Td, ThetaT)
GRT.RFb.fun(Temp, Tc, Tb, Td, ThetaT)
```

Arguments

The 'GR.funName()' functions have no arguments. The general purpose 'GR.funName.fun()' functions have some of the following arguments (depending on function):

Temp	Temperature variable
k	regression parameter
beta	regression parameter
Tc	ceiling temperature
Tb	base temperature
To	optimal temperature
Td	close-to-optimal temperature (Rowse-Fintch-Savage equation)
ThetaT	Hydro-time parameter

Details

All these functions are named according to this rule: 'GRT' (Germination Rate Temperature), followed by the 'function name' (e.g., BS, RF, M, Ex, YL). The R functions 'GR.funName().fun' are generic R functions, that are meant to be used for general purposes, such as plotting or predicting. The corresponding 'GR.funName()' (without the '.fun' ending) are meant to be used for fitting with the 'drm()' function, within the 'drc' package.

Value

The 'GRT.funName.fun()' functions return a vector of responses, for given values of temperature and parameters. The 'GRT.funName()' functions return a list containing the nonlinear function, the self starter function, the parameter names and other items which are internally used by the 'drm()' function.

Author(s)

Andrea Onofri

References

https://www.statforbiology.com/2023/stat_drcte_12-htt2step/

Examples

```
library(drcte)
Tval <- c(2, 5, 10, 15, 20, 25, 30, 35, 40)
GR <- c(0, 0, 0.209, 0.435, 0.759, 0.821, 0.417, 0.145, 0)
modM <- drm(GR ~ Tval, fct = GRT.M())
plot(modM, log="", xlim = c(0, 40), ylim=c(0,1.2),
      legendPos = c(5, 1.0), xlab = "Temperature (°C)")
```

hordeum

Germination of Hordeum spontaneum at different temperatures and water potentials

Description

This dataset was obtained from a germination assay with four replicated Petri dishes with 20 seeds, tested at six different water potential levels (0, -0.3, -0.6, -0.9, -1.2 and -1.5 MPa). Osmotic potentials were produced using variable amount of polyethylene glycol (PEG, molecular weight 8000) adjusted for the temperature level. Petri dishes were incubated at six constant temperature levels (8, 12, 16, 20, 24 and 28 °C), under a photoperiod of 12 h. Germinated seeds (radicle protrusion > 3 mm) were counted and removed daily for 20 days.

Usage

```
data("hordeum")
```

Format

A data frame with 3024 observations on the following 8 variables.

temp a numeric vector: temperature level

water a numeric vector: water potential level

Dish a numeric vector: code for Petri dishes

timeBef a numeric vector: beginning of scoring interval

timeAf a numeric vector: end of scoring interval

nViable a numeric vector: number of viable seeds at the beginning of assay, in each dish

nSeeds a numeric vector: number of germinated seeds, between timeBef and timeAf

nCum a numeric vector: cumulative number of germinated seeds at timeAf

Details

This dataset was analysed in the time-to-event framework in Onofri et al (2018). See Example 2.

Source

Mesgaran, MB, A Onofri, HR Mashhadi, RD Cousens (2017) Water availability shifts the optimal temperatures for seed germination: A modelling approach. *Ecological Modelling* 351:87–95

References

Mesgaran, MB, A Onofri, HR Mashhadi, RD Cousens (2017) Water availability shifts the optimal temperatures for seed germination: A modelling approach. *Ecological Modelling* 351:87–95

Onofri, A, P Benincasa, MB Mesgaran, C Ritz (2018) Hydrothermal-time-to-event models for seed germination. *European Journal of Agronomy* 101:129–139

Examples

```
# Fitting a hydrotime model
data(rape)
gmod <- drmtc(nSeeds ~ timeBef + timeAf + Psi,
             fct=HTnorm(), data=rape)
summary(gmod)

jackGroupSE(gmod, rape, rape$Dish)
```

HTE1 *Hydrotime model with log-logistic distribution of germination time (Onofri et al., 2018)*

Description

This model relates the time-course of the proportion of germinated seeds to the water potential in the substrate. It is based on a truncated log-logistic distribution of germination time:

$$P(t) = \frac{d}{1 + \exp[b(\log(x) - \log(e))]}$$

where two of the three usual parameters ('d' and 'e') are expressed as functions of water potential (Ψ). In this function, the two submodels are: (1) for the parameter 'd', we used a shifted exponential function:

$$d = G \left[1 - \exp \left(\frac{\Psi - \Psi_b}{\sigma_{\Psi_b}} \right) \right]$$

while, (2) for the parameter 'e' we considered that its inverse corresponds to the median Germination Rate within the population (i.e. $1/e = GR_{50}$) and modelled this latter parameter as:

$$GR_{50} = \frac{\Psi - \Psi_b}{\theta_H}$$

The 'HTE1.fun()' is a generic function, which can be used for plotting or other applications, while the 'HTE1()' function is meant to be used for model fitting with the 'drmte()' function in the 'drcte()' package.

Usage

```
HTE1(fixed = c(NA, NA, NA, NA, NA),
     names = c("G", "Psi_b", "sigmaPsi_b", "thetaH", "b"))
HTE1.fun(time, Psi, G, Psi_b, sigmaPsi_b, thetaH, b)
```

Arguments

These functions have the following arguments:

fixed	numeric vector. Specifies which parameters are fixed and at what value they are fixed. NAs for parameter that are not fixed. At the moment, only the parameter G can be held fixed.
names	a vector of character strings giving the names of the parameters. The default is reasonable.
time	time
Psi	water potential in the substrate
G	maximum germination capability of the seed lot

Psib	base water potential
sigmaPsib	standard deviation for the base water potential within the seed lot
thetaH	hydro-time constant
b	shape parameter for the cumulative distribution function of germination time

Details

The detail of this time-to-event model and the meaning of parameters are described in Onofri et al. (2018). See Table 2, where 'HTE1()' is abbreviated as HTE.

Value

The 'HTE1.fun()' function returns the proportion of germinated seeds, for any given values of time and water potential in the substrate. The 'HTE1()' function returns a list containing the nonlinear function, the self-starter function, the parameter names and other items which are internally used by the 'drmtte()' function.

Author(s)

Andrea Onofri

References

Onofri, A., Benincasa, P., Mesgaran, M.B., Ritz, C., 2018. Hydrothermal-time-to-event models for seed germination. *European Journal of Agronomy* 101, 129–139. https://www.statforbiology.com/2020/stat_seedgermination_ht1step/

Examples

```
data(rape)
modHTE <- drmtte( nSeeds ~ timeBef + timeAf + Psi,
                 data=rape, fct=HTE1())
summary(modHTE)
```

HTE2	<i>Hydrotime model with log-logistic distribution of germination time (Onofri et al., 2018)</i>
------	---

Description

This model relates the time-course of the proportion of germinated seeds to the water potential in the substrate. It is based on a truncated log-logistic distribution of germination time:

$$P(t) = \frac{d}{1 + \exp[b(\log(x) - \log(e))]}$$

where two of the three usual parameters ('d' and 'e') are expressed as functions of water potential (Ψ). In this function, the two submodels are: (1) for the parameter 'd', we used a shifted exponential function:

$$d = G \left[1 - \exp \left(\frac{\Psi - \Psi_b}{\sigma_{\Psi_b}} \right) \right]$$

while, (2) for the parameter 'e' we considered that its inverse corresponds to the median Germination Rate within the population (i.e. $1/e = GR_{50}$) and modelled this latter parameter as:

$$GR_{50} = \frac{[\Psi - \Psi_b]^2}{\theta_H}$$

The difference with the 'HTE1()' function is that, in this case, the relationship between GR50 and water potential is not linear, but curvilinear (convex down). The 'HTE2.fun()' is a generic function, which can be used for plotting or other applications, while the 'HTE2()' function is meant to be used for model fitting with the 'drmtc()' function in the 'drctc()' package.

Usage

```
HTE2(fixed = c(NA, NA, NA, NA, NA),
     names = c("G", "Psib", "sigmaPsib", "thetaH", "b"))
HTE2.fun(time, Psi, G, Psib, sigmaPsib, thetaH, b)
```

Arguments

These functions have the following arguments:

fixed	numeric vector. Specifies which parameters are fixed and at what value they are fixed. NAs for parameter that are not fixed. At the moment, only the parameter G can be held fixed.
names	a vector of character strings giving the names of the parameters. The default is reasonable.
time	time
Psi	water potential in the substrate
G	maximum germination capability of the seed lot
Psib	base water potential
sigmaPsib	standard deviation for the base water potential within the seed lot
thetaH	hydro-time constant
b	shape parameter for the cumulative distribution function of germination time

Details

No more detail, at the moment.

Value

The 'HTE2.fun()' function returns the proportion of germinated seeds, for any given values of time and water potential in the substrate, depending on model parameters G, Psib, sigmaPsib, thetaH and b. The 'HTE2()' function returns a list containing the nonlinear function, the self starter function, the parameter names and other items which are internally used by the 'drmtc()' function.

Author(s)

Andrea Onofri

References

Onofri, A., Benincasa, P., Mesgaran, M.B., Ritz, C., 2018. Hydrothermal-time-to-event models for seed germination. *European Journal of Agronomy* 101, 129–139. https://www.statforbiology.com/2020/stat_seedgermination_ht1step/

Examples

```
# Fitting model
data(rape)
modHTE2 <- drmtc( nSeeds ~ timeBef + timeAf + Psi,
                 data=rape, fct=HTE2())
summary(modHTE2)
```

HTE3

Hydrotime model with log-logistic distribution of germination time (Onofri et al., 2018)

Description

This model relates the time-course of the proportion of germinated seeds to the water potential in the substrate. It is based on a truncated log-logistic distribution of germination time:

$$P(t) = \frac{d}{1 + \exp[b(\log(x) - \log(e))]}$$

where two of the three usual parameters ('d' and 'e') are expressed as functions of water potential (Ψ). In this function, the two submodels are: (1) for the parameter 'd', we used a shifted exponential function:

$$d = G \left[1 - \exp \left(\frac{\Psi - \Psi_b}{\sigma_{\Psi_b}} \right) \right]$$

while, (2) for the parameter 'e' we considered that its inverse corresponds to the median Germination Rate within the population (i.e. $1/e = GR_{50}$) and modelled this latter parameter as:

$$GR_{50} = \frac{\Psi^2 - \Psi_b^2}{\theta_H}$$

The difference with the 'HTE1()' function is that, in this case, the relationship between GR50 and water potential is not linear, but curvilinear (convex up). The 'HTE3.fun()' is a generic function, which can be used for plotting or other applications, while the 'HTE3()' function is meant to be used for model fitting with the 'drmte()' function in the 'drcte()' package.

Usage

```
HTE3(fixed = c(NA, NA, NA, NA, NA),
      names = c("G", "Psib", "sigmaPsib", "thetaH", "b"))
HTE3.fun(time, Psi, G, Psib, sigmaPsib, thetaH, b)
```

Arguments

These functions have the following arguments:

fixed	numeric vector. Specifies which parameters are fixed and at what value they are fixed. NAs for parameter that are not fixed. At the moment, only the parameter G can be held fixed.
names	a vector of character strings giving the names of the parameters. The default is reasonable.
time	time
Psi	water potential in the substrate
G	maximum germination capability of the seed lot
Psib	base water potential
sigmaPsib	standard deviation for the base water potential within the seed lot
thetaH	hydro-time constant
b	shape parameter for the cumulative distribution function of germination time

Details

No more detail, at the moment.

Value

The 'HTE3.fun()' function returns the proportion of germinated seeds, for any given values of time and water potential in the substrate, depending on model parameters G, Psib, sigmaPsib, thetaH and b. The 'HTE3()' function returns a list containing the nonlinear function, the self starter function, the parameter names and other items which are internally used by the 'drmte()' function.

Author(s)

Andrea Onofri

References

Onofri, A., Benincasa, P., Mesgaran, M.B., Ritz, C., 2018. Hydrothermal-time-to-event models for seed germination. *European Journal of Agronomy* 101, 129–139. https://www.statforbiology.com/2020/stat_seedgermination_ht1step/

Examples

```
# Fitting model
data(rape)
modHTE3 <- drmte( nSeeds ~ timeBef + timeAf + Psi,
                 data = rape, fct=HTE3())
summary(modHTE3)
```

HTex

Hydrotime model with Type II Extreme Value distribution of base water potential (Mesgaran et al., 2013)

Description

This model relates the time-course of the proportion of germinated seeds to the water potential in the substrate. It is based on Type II Extreme Value distribution of base water potential within the seed lot. The equation is:

$$P(t) = 1 - \exp \left\{ - \exp \left[\frac{\Psi - (\theta_H/t) - \mu}{\sigma} \right] \right\}$$

In contrast to other hydrotime models (e.g., 'HTE1()', 'HTE2()', 'HTE3()'), describing the distribution of germination time), this models describes the distribution of base water potential within the population, while the distribution of germination times is indirectly modelled, but it is not, in itself, gaussian (you see that 't' is at the denominator). The 'HTex.fun()' is a generic function, which can be used for plotting or other applications, while the 'HTex()' function is meant to be used for model fitting with the 'drmte()' function in the 'drcte()' package.

Usage

```
HTex()
HTex.fun(time, Psi, thetaH, mu, sigma)
```

Arguments

The 'HTex()' function has no arguments. The 'HTex.fun()' function has the following arguments:

time	time
Psi	water potential in the substrate
thetaH	hydro-time constant
mu	location parameter for the Type II Extreme Value distribution
sigma	scale parameter for the Type II Extreme Value distribution

Details

The detail of this time-to-event model and the meaning of parameters are described in Mesgaran et al. (2013).

Value

The 'HTex.fun()' function returns the proportion of germinated seeds, for any given values of time and water potential in the substrate. The 'HTex()' function returns a list containing the nonlinear function, the self-starter function, the parameter names and other items which are internally used by the 'drmtc()' function.

Note

This function is for use with the function 'drmtc()'

Author(s)

Andrea Onofri

References

Mesgaran, M.B., Mashhadi, H.R., Alizadeh, H., Hunt, J., Young, K.R., Cousens, R.D., 2013. Importance of distribution function selection for hydrothermal time models of seed germination. *Weed Research* 53, 89–101.

Examples

```
data(rape)
modg <- drm( propCum ~ timeAf + Psi, fct=HTex(), data=rape)
summary(modg)
```

HTG

*Hydrotime model with Gumbel distribution of base water potential
(Mesgaran et al., 2013)*

Description

This model relates the time-course of the proportion of germinated seeds to the water potential in the substrate. It is based on a Gumbel distribution of base water potential within the seed lot. The equation is:

$$P(t) = \exp \left\{ - \exp \left[- \left(\frac{\Psi - (\theta_H/t) - \mu}{\sigma} \right) \right] \right\}$$

In contrast to other hydrotime models (e.g., 'HTE1()', 'HTE2()', 'HTE3()'), describing the distribution of germination time), this models describes the distribution of base water potential within the population, while the distribution of germination times is indirectly modelled, but it is not, in itself, gaussian (you see that 't' is at the denominator). The 'HTG.fun()' is a generic function, which can be used for plotting or other applications, while the 'HTG()' function is meant to be used for model fitting with the 'drmtc()' function in the 'drctc()' package.

Usage

```
HTG()  
HTG.fun(time, Psi, ThetaH, mu, sigma)
```

Arguments

The 'HTG()' function has no arguments. The 'HTG.fun()' function has the following arguments:

time	time
Psi	water potential in the substrate
ThetaH	hydro-time constant
mu	location parameter for the Gumbel distribution
sigma	scale parameter for the Gumbel distribution

Details

The detail of this time-to-event model and the meaning of parameters are described in Mesgaran et al. (2013).

Value

The 'HTG.fun()' function returns the proportion of germinated seeds, for any given values of time and water potential in the substrate. The 'HTG()' function returns a list containing the nonlinear function, the self-starter function, the parameter names and other items which are internally used by the 'drmtc()' function.

Note

This function is for use with the function 'drmtc()'

Author(s)

Andrea Onofri

References

Mesgaran, M.B., Mashhadi, H.R., Alizadeh, H., Hunt, J., Young, K.R., Cousens, R.D., 2013. Importance of distribution function selection for hydrothermal time models of seed germination. *Weed Research* 53, 89–101.

Examples

```
data(rape)  
modg <- drm( propCum ~ timeAf + Psi, fct=HTG(), data=rape)  
summary(modg)
```

HTL *Hydrotime model with logistic distribution of base water potential*
(Mesgaran et al., 2013)

Description

This model relates the time-course of the proportion of germinated seeds to the water potential in the substrate. It is based on a logistic distribution of base water potential within the seed lot. The equation is:

$$P(t) = \frac{1}{1 + \exp\left[-\frac{\Psi - (\theta_H/t) - \Psi_b}{\sigma}\right]}$$

In contrast to other hydrotime models (e.g., 'HTE1()', 'HTE2()', 'HTE3()'), describing the distribution of germination time), this models describes the distribution of base water potential within the population, while the distribution of germination times is indirectly modelled, but it is not, in itself, gaussian (you see that 't' is at the denominator). The 'HTL.fun()' is a generic function, which can be used for plotting or other applications, while the 'HTL()' function is meant to be used for model fitting with the 'drmtc()' function in the 'drcte()' package.

Usage

```
HTL()
HTL.fun(time, Psi, ThetaH, Psib50, sigma)
```

Arguments

The 'HTL()' function has no arguments. The 'HTL.fun()' function has the following arguments:

time	time
Psi	water potential in the substrate
ThetaH	hydro-time constant
Psib50	median base water potential
sigma	scale parameter for the logistic distribution

Details

The detail of this time-to-event model and the meaning of parameters are described in Mesgaran et al. (2013).

Value

The 'HTL.fun()' function returns the proportion of germinated seeds, for any given values of time and water potential in the substrate. The 'HTL()' function returns a list containing the nonlinear function, the self-starter function, the parameter names and other items which are internally used by the 'drmtc()' function.

Note

This function is for use with the function 'drmtc()'

Author(s)

Andrea Onofri

References

Mesgaran, M.B., Mashhadi, H.R., Alizadeh, H., Hunt, J., Young, K.R., Cousens, R.D., 2013. Importance of distribution function selection for hydrothermal time models of seed germination. *Weed Research* 53, 89–101.

Examples

```
data(rape)
modg <- drm( propCum ~ timeAf + Psi, fct=HTL(), data=rape)
summary(modg)
```

HTLL

Hydrotime model with log-logistic distribution of base water potential (Mesgaran et al., 2013)

Description

This model relates the time-course of the proportion of germinated seeds to the water potential in the substrate. It is based on a log-logistic distribution of base water potential within the seed lot. The equation is:

$$P(t) = \frac{1}{1 + \exp \left\{ \frac{\log \left[\Psi - \left(\frac{\theta H}{t} \right) + \delta \right] - \log(\Psi_b + \delta)}{\sigma} \right\}}$$

In contrast to other hydrotime models (e.g., 'HTE1()', 'HTE2()', 'HTE3()'), describing the distribution of germination time), this models describes the distribution of base water potential within the population, while the distribution of germination times is indirectly modelled, but it is not, in itself, gaussian (you see that 't' is at the denominator). The 'HTLL.fun()' is a generic function, which can be used for plotting or other applications, while the 'HTLL()' function is meant to be used for model fitting with the 'drmtc()' function in the 'drctc()' package.

Usage

```
HTLL()
HTLL.fun(time, Psi, thetaH, delta, Psib50, sigma)
```

Arguments

The 'HTLL()' function has no arguments. The 'HTLL.fun()' function has the following arguments:

time	time
Psi	water potential in the substrate
thetaH	hydro-time constant
delta	shifting parameter to allow for negative water potential levels
Psi _{b50}	median base water potential
sigma	scale parameter for the log-logistic distribution

Details

The detail of this time-to-event model and the meaning of parameters are described in Mesgaran et al. (2013).

Value

The 'HTLL.fun()' function returns the proportion of germinated seeds, for any given values of time and water potential in the substrate. The 'HTLL()' function returns a list containing the nonlinear function, the self-starter function, the parameter names and other items which are internally used by the 'drmte()' function.

Note

This function is for use with the function 'drmte()'

Author(s)

Andrea Onofri

References

Mesgaran, M.B., Mashhadi, H.R., Alizadeh, H., Hunt, J., Young, K.R., Cousens, R.D., 2013. Importance of distribution function selection for hydrothermal time models of seed germination. *Weed Research* 53, 89–101.

Examples

```
data(rape)
modg <- drm( propCum ~ timeAf + Psi, fct=HTLL(), data=rape)
summary(modg)
```

HTnorm	<i>Hydrotime model with normal distribution of base water potential (Bradford, 2002)</i>
--------	--

Description

This model relates the time-course of the proportion of germinated seeds to the water potential in the substrate. It is based on a normal distribution of base water potential within the seed lot. The equation is:

$$P(t) = \Phi \left\{ \frac{\Psi - (\theta_H/t) - \Psi_b}{\sigma_{\Psi_b}} \right\}$$

where Φ is a gaussian cumulative distribution function for base water potential. In contrast to other hydrotime models (e.g., 'HTE1()', 'HTE2()', 'HTE3()'), describing the distribution of germination time), this models describes the distribution of base water potential within the population, while the distribution of germination times is indirectly modelled, but it is not, in itself, gaussian (you see that 't' is at the denominator). The 'HTNorm.fun()' is a generic function, which can be used for plotting or other applications, while the 'HTNorm()' function is meant to be used for model fitting with the 'drmtc()' function in the 'drctc()' package.

Usage

```
HTnorm()
HTnorm.fun(time, Psi, ThetaH, Psib50, sigmaPsib)
```

Arguments

The 'HTNorm()' function has no arguments. The 'HTNorm.fun()' function has the following arguments:

time	time
Psi	water potential in the substrate
ThetaH	hydro-time constant
Psib50	median base water potential
sigmaPsib	standard deviation for the base water potential within the seed lot

Details

The detail of this time-to-event model and the meaning of parameters are described in Bradford (2002).

Value

The 'HTNorm.fun()' function returns the proportion of germinated seeds, for any given values of time and water potential in the substrate. The 'HTNorm()' function returns a list containing the nonlinear function, the self-starter function, the parameter names and other items which are internally used by the 'drmtc()' function.

Note

This function is for use with the R function 'drmtc()'

Author(s)

Andrea Onofri

References

Bradford, K.J., 2002. Applications of hydrothermal time to quantifying and modeling seed germination and dormancy. *Weed Science* 50, 248–260.

Examples

```
data(rape)
modg <- drm( propCum ~ timeAf + Psi, fct=HTnorm(), data=rape)
summary(modg)
```

HTTEM

Hydro-thermal-time model with log-logistic distribution of germination time (Onofri et al., 2018)

Description

This model relates the time-course of the proportion of germinated seeds to the water potential and temperature in the substrate. It is based on a truncated log-logistic distribution of germination time:

$$P(t) = \frac{d}{1 + \exp[b(\log(x) - \log(e))]}$$

where two of the three usual parameters ('d' and 'e') are expressed as functions of water potential (Ψ) and temperature (T). In the function 'HTTEM()', we implemented the following submodels: (1) for the parameter 'd', we implemented a shifted exponential function:

$$d = G \left[1 - \exp\left(\frac{\Psi - \Psi_b - k(T - T_b)}{\sigma_{\Psi_b}}\right) \right]$$

while, (2) for the parameter 'e' we considered that its inverse corresponds to the median Germination Rate within the population (i.e. $1/e = GR_{50}$) and modelled this latter parameter as:

$$GR_{50} = \frac{T - T_b}{\theta_{HT}} [\Psi - \Psi_b - k(T - T_b)]$$

The 'HTTEM.fun()' is a generic function, which can be used for plotting or other applications, while the 'HTTEM()' function is meant to be used for model fitting with the 'drmtc()' function in the 'drctc()' package.

Usage

```
HTTEM()
HTTEM.fun(time, Psi, Temp, G, Psib, kt, Tb, sigmaPsib, ThetaHT, b)
```

Arguments

The 'HTTEM' function has no arguments. The 'HTTEM.fun()' has the following arguments:

time	time
Psi	water potential in the substrate
Temp	temperature
G	maximum germination capability of the seed lot
Psib	base water potential
kt	parameter measuring the effect of temperature on base water potential
Tb	base temperature
sigmaPsib	standard deviation for the base water potential within the seed lot
ThetaHT	hydro-thermal-time parameter
b	shape parameter for the cumulative distribution function of germination time

Details

The detail of this time-to-event model and the meaning of parameters are described in Onofri et al. (2018). See Table 2, where 'HTTEM()' is abbreviated as HTTE.

Value

The 'HTTEM.fun()' function returns the proportion of germinated seeds, for any given values of time and water potential in the substrate. The 'HTTEM()' function returns a list containing the nonlinear function, the parameter names and other items which are internally used by the 'drcte()' function. At the moment, there is no self-starting function and starting parameters for fitting must be provided within the 'drcte' function.

Author(s)

Andrea Onofri

References

Onofri, A., Benincasa, P., Mesgaran, M.B., Ritz, C., 2018. Hydrothermal-time-to-event models for seed germination. *European Journal of Agronomy* 101, 129–139. https://www.statforbiology.com/2023/stat_drcte_10-examplehtte/

Examples

```
data(hordeum)

modHTTEM <- drmtc(nSeeds ~ timeBef + timeAf + water + temp,
                 data=hordeum,
                 fct = HTTEM(),
                 start=c(0.8,-2, 0.05, 3, 0.2, 2000, 0.5))
summary(modHTTEM)
```

HTTLL	<i>Hydro-thermal-time model with log-logistic distribution of base water potential (Mesgaran et al., 2017)</i>
-------	--

Description

This model relates the time-course of the proportion of germinated seeds to the water potential and temperature in the substrate and it is based on a log-logistic distribution of base water potential within the seed lot. Two similar functions are available within the 'drcSeedGerm' package: the first one is 'HTTLL.M()' that assumes that the base water potential decreases with temperature for any $T > T_b$. The equation is:

$$P(t, T, \Psi) = \frac{1}{1 + \exp \left\{ - \frac{\log \left[\Psi - \left(\frac{\theta_{HT}}{t(T - T_b)} \right) + \delta \right] - \log[\Psi_b + K_t(T - T_b) + \delta]}{\sigma_{\Psi_b}} \right\}}$$

The second function is 'HTTLL.BS()' and it assumes that the base water potential decreases with temperature only for any $T > T_o$, where T_o is the optimal temperature level. For this case, the element $K_t(T - T_b)$ is modified as $K_t[\max(T, T_o) - T_o]$ and the optimal temperature T_o is included as an explicit parameter.

The 'HTTLL.M.fun()' and 'HTTLL.BS.fun()' are two generic functions, which can be used for plotting or other applications, while the 'HTTLL.M()' and 'HTTLL.BS()' functions are meant to be used for model fitting with the 'drmtc()' function in the 'drcTe()' package.

Usage

```
HTTLL.M()
HTTLL.BS()
HTTLL.M.fun(time, Psi, Temp, thetaHT, Tb, Psib50, Kt, delta, sigmaPsib)
HTTLL.BS.fun(time, Psi, Temp, thetaHT, Tb, To, Psib50, Kt, delta, sigmaPsib)
```

Arguments

The 'HTTLL.M()' and 'HTTLL.BS()' functions have no arguments. The 'HTTLL.M.fun()' and the 'HTTLL.BS.fun()' functions have the following arguments:

```
time           time
```

Psi	water potential in the substrate
Temp	temperature
thetaHT	hydro-time parameter
Tb	base temperature
To	optimal temperature
Psib50	median base water potential
Kt	parameter measuring the effect of temperature on base water potential
delta	shifting parameter to allow for a log-logistic distribution of negative values for base water potential
sigmaPsib	scale parameter for log-logistic distribution

Details

The detail of this time-to-event model and the meaning of parameters are described in Mesgaran et al. (2017).

Value

The 'HTTLL.M.fun()' and 'HTTLL.BS.fun()' functions return the proportion of germinated seeds, for any given values of time, water potential and temperature in the substrate. The 'HTTLL.M()' and 'HTTLL.BS()' functions return a list containing the nonlinear function, the self-starter function, the parameter names and other items which are internally used by the 'drmtc()' function.

Note

This function is for use with the function 'drmtc()'

Author(s)

Andrea Onofri

References

Bradford, K.J., 2002. Applications of hydrothermal time to quantifying and modeling seed germination and dormancy. *Weed Science* 50, 248–260.

Examples

```
data(hordeum)

modHTTLL.M <- drmtc(nSeeds ~ timeBef + timeAf + water + temp,
                  data=hordeum,
                  fct = HTTLL.M(),
                  start=c(832,-2.5, -3, 0.07, 3, 0.5))
summary(modHTTLL.M)

modHTTLL.BS <- drmtc(nSeeds ~ timeBef + timeAf + water + temp,
                   data=hordeum,
```

```
fct = HTLL.BS(),
start=c(932,-2.5, 15, -3, 0.07, 3, 0.5))
summary(modHTLL.BS)
```

HTTnorm.M	<i>Hydro-thermal-time model with normal distribution of base water potential (Bradford, 2002)</i>
-----------	---

Description

This model relates the time-course of the proportion of germinated seeds to the water potential and temperature in the substrate and it is based on a normal distribution of base water potential within the seed lot. Two similar functions are available within the 'drcSeedGerm' package: the first one is 'HTTnorm.M()' that assumes that the base water potential decreases with temperature for any $T > T_b$. The equation is:

$$P(t, T, \Psi) = \Phi \left\{ \frac{\Psi - \left[\frac{\theta_{HT}}{t(T - T_b)} \right] - [\Psi_b + K_t(T - T_b)]}{\sigma_{\Psi_b}} \right\}$$

where Φ is a gaussian cumulative distribution function for base water potential.

The second function is 'HTTNorm.BS()' and it assumes that the base water potential decreases with temperature only for any $T > T_o$, where T_o is the optimal temperature level. For this case, the element $K_t(T - T_b)$ is modified as $K_t[\max(T, T_o) - T_o]$ and the optimal temperature T_o is included as an explicit parameter.

The 'HTTNorm.M.fun()' and 'HTTNorm.BS.fun()' are two generic functions, which can be used for plotting or other applications, while the 'HTTnorm.M()' and 'HTTnorm.BS()' functions are meant to be used for model fitting with the 'drmte()' function in the 'drcTe()' package.

Usage

```
HTTnorm.M()
HTTnorm.BS()
HTTnorm.M.fun(time, Psi, Temp, thetaHT, Tb, Psib50, Kt, sigmaPsib)
HTTnorm.BS.fun(time, Psi, Temp, thetaHT, Tb, To, Psib50, Kt, sigmaPsib)
```

Arguments

The 'HTTnorm.M()' and 'HTTnorm.BS()' functions have no arguments. The 'HTTnorm.M.fun()' and the 'HTTnorm.BS.fun()' functions have the following arguments:

time	time
Psi	water potential in the substrate
Temp	temperature
thetaHT	hydro-time parameter

Tb	base temperature
To	optimal temperature
Psib50	median base water potential
Kt	parameter measuring the effect of temperature on base water potential
sigmaPsib	standard deviation for the base water potential within the seed lot

Details

The detail of this time-to-event model and the meaning of parameters are described in Bradford (2002).

Value

The 'HTTnorm.M.fun()' and 'HTTnorm.BS.fun()' functions return the proportion of germinated seeds, for any given values of time, water potential and temperature in the substrate. The 'HTTnorm.M()' and 'HTTnorm.BS()' functions return a list containing the nonlinear function, the self-starter function, the parameter names and other items which are internally used by the 'drmtc()' function.

Note

This function is for use with the function 'drmtc()'

Author(s)

Andrea Onofri

References

Bradford, K.J., 2002. Applications of hydrothermal time to quantifying and modeling seed germination and dormancy. *Weed Science* 50, 248–260.

Examples

```
data(hordeum)

modHTTnorm.M <- drmtc(nSeeds ~ timeBef + timeAf + water + temp,
                    data=hordeum,
                    fct = HTTnorm.M(),
                    start=c(932,-2.5, -3, 0.07, 0.5))
summary(modHTTnorm.M)
```

HTW1

Hydrotime model with Weibull Type I distribution of base water potential (Mesgaran et al., 2013)

Description

This model relates the time-course of the proportion of germinated seeds to the water potential in the substrate. It is based on a Weibull Type I distribution of base water potential within the seed lot. The equation is:

$$P(t) = \exp \left\{ - \exp \left[- \frac{\log [\Psi - (\frac{\theta_H}{t}) + \delta] - \log(\Psi_b + \delta)}{\sigma} \right] \right\}$$

In contrast to other hydrotime models (e.g., 'HTE1()', 'HTE2()', 'HTE3()'), describing the distribution of germination time), this models describes the distribution of base water potential within the population, while the distribution of germination times is indirectly modelled, but it is not, in itself, gaussian (you see that 't' is at the denominator). The 'HTW1.fun()' is a generic function, which can be used for plotting or other applications, while the 'HTW1()' function is meant to be used for model fitting with the 'drnte()' function in the 'drcte()' package.

Usage

```
HTW1()
HTW1.fun(time, Psi, thetaH, delta, mu, sigma)
```

Arguments

The 'HTW1()' function has no arguments. The 'HTW1.fun()' function has the following arguments:

time	time
Psi	water potential in the substrate
thetaH	hydro-time constant
delta	shifting parameter to allow for negative water potential levels
mu	location parameter for the Type I Weibull distribution
sigma	scale parameter for the Type I Weibull Value distribution

Details

The detail of this time-to-event model and the meaning of parameters are described in Mesgaran et al. (2013).

Value

The 'HTW1.fun()' function returns the proportion of germinated seeds, for any given values of time and water potential in the substrate. The 'HTW1()' function returns a list containing the nonlinear function, the self-starter function, the parameter names and other items which are internally used by the 'drnte()' function.

Note

This function is for use with the function 'drmtc()'

Author(s)

Andrea Onofri

References

Mesgaran, M.B., Mashhadi, H.R., Alizadeh, H., Hunt, J., Young, K.R., Cousens, R.D., 2013. Importance of distribution function selection for hydrothermal time models of seed germination. *Weed Research* 53, 89–101.

Examples

```
data(rape)
modg <- drm( propCum ~ timeAf + Psi, fct=HTW1(), data=rape)
summary(modg)
```

HTW2

Hydrotime model with Weibull Type II distribution of base water potential (Mesgaran et al., 2013)

Description

This model relates the time-course of the proportion of germinated seeds to the water potential in the substrate. It is based on a Weibull Type II distribution of base water potential within the seed lot. The equation is:

$$P(t) = 1 - \exp \left\{ - \exp \left[\frac{\log \left[\Psi - \left(\frac{\theta_H}{t} \right) + \delta \right] - \log(\Psi_b + \delta)}{\sigma} \right] \right\}$$

In contrast to other hydrotime models (e.g., 'HTE1()', 'HTE2()', 'HTE3()'), describing the distribution of germination time), this models describes the distribution of base water potential within the population, while the distribution of germination times is indirectly modelled, but it is not, in itself, gaussian (you see that 't' is at the denominator). The 'HTW2.fun()' is a generic function, which can be used for plotting or other applications, while the 'HTW2()' function is meant to be used for model fitting with the 'drmtc()' function in the 'drctc()' package.

Usage

```
HTW2()
HTW2.fun(time, Psi, thetaH, delta, mu, sigma)
```

Arguments

The 'HTW2()' function has no arguments. The 'HTW2.fun()' function has the following arguments:

time	time
Psi	water potential in the substrate
thetaH	hydro-time constant
delta	shifting parameter to allow for negative water potential levels
mu	location parameter for the Type II Weibull distribution
sigma	scale parameter for the Type II Weibull Value distribution

Details

The detail of this time-to-event model and the meaning of parameters are described in Mesgaran et al. (2013).

Value

The 'HTW2.fun()' function returns the proportion of germinated seeds, for any given values of time and water potential in the substrate. The 'HTW2()' function returns a list containing the nonlinear function, the self-starter function, the parameter names and other items which are internally used by the 'drmtc()' function.

Note

This function is for use with the function 'drmtc()'

Author(s)

Andrea Onofri

References

Mesgaran, M.B., Mashhadi, H.R., Alizadeh, H., Hunt, J., Young, K.R., Cousens, R.D., 2013. Importance of distribution function selection for hydrothermal time models of seed germination. *Weed Research* 53, 89–101.

Examples

```
data(rape)
modg <- drm( propCum ~ timeAf + Psi, fct=HTW2(), data=rape)
summary(modg)
```

`jackGroupSE`*Delete-a-Petri-dish jackknife*

Description

This function estimates standard errors for regression model parameters by using the fully-iterated delete-a-group jackknife (Yu and Peng, 2008). This is asymptotically equivalent to the cluster-robust 'sandwich' variance estimator (Lipsitz, 1994).

Usage

```
jackGroupSE(obj, data, units)
```

Arguments

<code>obj</code>	a fitted model object
<code>data</code>	dataset containing the variables used in <code>obj</code>
<code>units</code>	a vector containing the code for Petri dishes

Details

Care should be taken to ensure that the assumptions for using a fully-iterated jackknife are valid.

Value

This function returns a data.frame containing the estimated parameters, the SEs and the robust SEs

Author(s)

Andrea Onofri

References

Yu, B., Peng, Y., 2008. Mixture cure models for multivariate survival data. *Computational Statistics and Data Analysis* 52, 1524–1532. Lipsitz, S.R., Dear, K.B.G., Zhao, L., 1994. Jackknife Estimators of Variance for Parameter Estimates from Estimating Equations with Applications to Clustered Survival Data. *Biometrics* 50, 842–846.

Examples

```
data(rape)

modHTE <- drmte( nSeeds ~ timeBef + timeAf + Psi,
                data=rape, fct=HTE1())
robust <- jackGroupSE(modHTE, rape, rape$Dish)
robust
```

makeDrm	<i>Reshape a seed germination datasets for time-to-event analyses (deprecated)</i>
---------	--

Description

MakeDrm: this function reshapes a common field book for germination assays into the form that is required for time-to-event analyses with `drm()` in the `drc` package and `drmte()` in the `drcte` package. The common field book has one row per each Petri dish and the counts of germinated seeds at each assessment time are listed in different columns.

Usage

```
makeDrm(counts, treat, nViable, moniTimes)
```

Arguments

<code>counts</code>	Dataframe listing the counts of germinated seeds in each Petri dish (rows) at each assessment time (columns).
<code>treat</code>	Dataframe listing for each dish the levels of each treatment. It has as many columns as there are treatments in the experiment.
<code>nViable</code>	A vector with the number of viable seeds per dish, at the beginning of the assay
<code>moniTimes</code>	Vector of monitoring times. It needs to be of the same length as the number of columns in 'counts'.

Value

This function returns a dataframe

Author(s)

Andrea Onofri

makeDrm2	<i>Reshape a seed germination datasets for time-to-event model fitting.</i>
----------	---

Description

This function reshapes a datasets organised as necessary for nonlinear regression into the kind of dataset required by the `drmte()` function in the 'drcte' package. It works with either the counts of germinated seeds at each monitoring time or the cumulative counts at each monitoring time.

Usage

```
makeDrm2(counts, treat, nViable, moniTimes, Dish, cumulative = TRUE)
```

Arguments

counts	Vector listing the counts of germinated seeds in each Petri dish at each assessment time.
treat	Dataframe listing, for each row of data, the corresponding level of experimental factors (one factor per column)
nViable	A vector listing the number of viable seeds, at the beginning of the assay. This number is the same for all observations belonging to the same dish.
moniTimes	Vector of monitoring times.
Dish	Vector of codes for dishes.
cumulative	Logical: True if counts are cumulative, False if they are not.

Value

Returns a dataframe

Author(s)

Andrea Onofri

Examples

```
# makeDrm2 (deprecated)
data(lotusCum)
moniTime <- lotusCum$Time
count <- lotusCum$nCum
nViable <- rep(25, length(lotusCum[,1]))
Dish <- as.factor(lotusCum$Dish)
treatGroups <- lotusCum[,1]
dataset_sd <- makeDrm2(count, treatGroups, nViable, moniTime, Dish)
head(dataset_sd)
count <- lotusCum$nSeeds
dataset_sd <- makeDrm2(count, treatGroups, nViable, moniTime, Dish, cumulative = FALSE)
head(dataset_sd)
```

phalaris

A series of germination assays with Phalaris minor

Description

The germination of *Phalaris minor* was assayed at 6 water potential levels and 6 temperature levels, on four replicated Petri dishes at each temperature and water potential. Twenty-five seeds per Petri dish were used. Inspections were made daily for twenty days.

Usage

```
data("phalaris")
```

Format

A data frame with 3024 observations on the following 9 variables.

`temp` a numeric vector with the temperature level

`water` a numeric vector with the water potential level

`Dish` a numeric vector with the coding for Petri dishes

`timeBef` a numeric vector, with the start time for each inspection interval

`timeAf` a numeric vector, with the end time for each inspection interval

`nViable` a numeric vector, with the number of viable seeds per dish, at the beginning of the assay

`nSeeds` a numeric vector, with the number of germinated seeds during each inspection interval

`nCum` a numeric vector, with the cumulative number of germinated seeds at each assessment time

`propCum` a numeric vector, the cumulative proportion of germinated seeds at each assessment time

Details

The variable 'timeAf' contains the value 'Inf' (Infinity), that corresponds to the seeds which did not germinate during the assay, for which the germination times might be comprised from the last assessment time to infinity.

Author(s)

Mohsen Mesgaran

Source

Mesgaran, MB, A Onofri, HR Mashhadi, RD Cousens (2017) Water availability shifts the optimal temperatures for seed germination: A modelling approach. *Ecological Modelling* 351, 87–95

References

Mesgaran, MB, A Onofri, HR Mashhadi, RD Cousens (2017) Water availability shifts the optimal temperatures for seed germination: A modelling approach. *Ecological Modelling* 351, 87–95

Examples

`data(phalaris)`

Pmax_models	<i>Effect of environmental water potential or temperature on the germination capability of a seed lot</i>
-------------	---

Description

These models are used to describe the germination capability of a seed lot, depending on the environmental water potential or temperature.

Usage

```
PmaxPsi1(fixed = c(NA, NA, NA), names = c("G", "Psib", "sigma"))
PmaxT1(fixed = c(NA, NA, NA), names = c("G", "Tc", "sigmaTc"))
PmaxPsi1.fun(Psi, G, Psib, sigma)
PmaxT1.fun(Temp, G, Tc, sigmaTc)
```

Arguments

fixed	numeric vector. Specifies which parameters are fixed and at what value they are fixed. Defaults to NAs for parameter that are not fixed.
names	a vector of character strings giving the names of the parameters (should not contain ":"). The default is reasonable.
Psi	Water potential level
Temp	Temperature level
G	Maximum proportion of germinated seeds
Psib	Base water potential level
sigma	Variability of base water potential within the seed lot
Tc	Ceiling temperature level
sigmaTc	Variability of ceiling temperature level within the population

Details

The R functions 'PmaxPsi1()' and 'PmaxT1()' are meant to be used for model fitting with the 'drm()' function, within the 'drc' package. On the other hand, 'PmaxPsi1.fun()' and 'PmaxT1.fun()' are general purpose functions, to be used for plotting or other applications.

Value

'PmaxPsi1()' and 'PmaxT1()' return a list containing the nonlinear function, the self starter function and the parameter names, that are internally used for model fitting. 'PmaxPsi1.fun()' and 'PmaxT1.fun()' return the maximum proportion of germinated seeds for any given level of temperature or water potential in the substrate.

Author(s)

Andrea Onofri

References

https://www.statforbiology.com/2023/stat_drcte_12-htt2step/

Examples

```
library(drcte)
# Pmax vs Psi (shifted exponential)
Psi <- seq(-2.2, 0, by = 0.2)
Pmax <- c(0, 0, 0.076, 0.413, 0.514, 0.643, 0.712,
          0.832, 0.865, 0.849, 0.89, 0.90)
mod <- drm(Pmax ~ Psi, fct = PmaxPsi1())
summary(mod)

# Pmax vs Psi (shifted exponential, with asymptote)
Psi <- seq(-2.2, 0, by = 0.2)
Pmax <- c(0, 0, 0.076, 0.413, 0.514, 0.643, 0.712,
          0.832, 0.865, 0.849, 0.89, 0.90)
mod <- drm(Pmax ~ Psi, fct = PmaxPsi1(fixed = c(1, NA, NA)))
summary(mod)

# Pmax vs temperature
Tval <- c(0, 2.5, 5, 7.5, 10, 12.5, 15, 17.5,
          20, 22.5, 25, 27.5, 30, 32.5, 35)
Pmax2 <- c(0.79, 0.81, 0.807, 0.776, 0.83,
           0.73, 0.744, 0.73, 0.828, 0.818,
           0.805, 0.706, 0.41, 0.002, 0)
mod2 <- drm(Pmax2 ~ Tval, fct = PmaxT1())
summary(mod2)
```

rape

Germination data from an assay of rapeseed at decreasing water potential levels

Description

This files describes the germination of rapeseed (cv. Excalibur) at different water potential levels in the substrate.

Usage

```
data("rape")
```

Format

A data frame with 294 observations on the following 7 variables.

Psi a numeric vector: waterpotential level

Dish a numeric vector: code for Petri dishes

timeBef a numeric vector: beginning of scoring interval

timeAf a numeric vector: end of scoring interval
 nSeeds a numeric vector: number of germinated seeds, between timeBef and timeAf
 nCum a numeric vector: cumulative number of germinated seeds at timeAf
 propCum a numeric vector: cumulative proportion of germinated seeds at timeAf

Source

Pace, R., Benincasa, P., Ghanem, M.E., Quinet, M., Lutts, S., 2012. Germination of untreated and primed seeds in rapeseed (*Brassica napus* var. *oleifera* Del.) under salinity and low matric potential. *Experimental Agriculture* 48, 238–251.

References

Pace, R., Benincasa, P., Ghanem, M.E., Quinet, M., Lutts, S., 2012. Germination of untreated and primed seeds in rapeseed (*Brassica napus* var. *oleifera* Del.) under salinity and low matric potential. *Experimental Agriculture* 48, 238–251.

Examples

```
#Fitting a hydrotime model
library(drcte)
data(rape)
gmod <- drmte(nSeeds ~ timeBef + timeAf + Psi,
              data = rape, fct = HTE1())
summary(gmod)
```

rape2G

Germination data from an assay of rapeseed at decreasing water potential levels

Description

This files describes the germination of rapeseed (cv. Excalibur and Toccata) at different water potential levels in the substrate.

Usage

```
data("rape")
```

Format

A data frame with 294 observations on the following 8 variables.

CV a factor: rape genotype
 Psi a numeric vector: waterpotential level
 Dish a numeric vector: code for Petri dishes
 timeBef a numeric vector: beginning of scoring interval

timeAf a numeric vector: end of scoring interval
 nSeeds a numeric vector: number of germinated seeds, between timeBef and timeAf
 nCum a numeric vector: cumulative number of germinated seeds at timeAf
 Prop a numeric vector: cumulative proportion of germinated seeds at timeAf

Source

Pace, R., Benincasa, P., Ghanem, M.E., Quinet, M., Lutts, S., 2012. Germination of untreated and primed seeds in rapeseed (*Brassica napus* var. *oleifera* Del.) under salinity and low matric potential. *Experimental Agriculture* 48, 238–251.

References

Pace, R., Benincasa, P., Ghanem, M.E., Quinet, M., Lutts, S., 2012. Germination of untreated and primed seeds in rapeseed (*Brassica napus* var. *oleifera* Del.) under salinity and low matric potential. *Experimental Agriculture* 48, 238–251.

Examples

```
#Fitting a hydrotime-to-event model
data(rape2G)
modHTE <- drmte( nSeeds ~ timeBef + timeAf + Psi,
                data=rape2G, fct=HTE1(),
                curveid = CV)
summary(modHTE)
```

Thermal-Time_models *Thermal-time models with log-logistic distribution of germination time (Onofri et al., 2018)*

Description

These models relates the time-course of germinations to the environmental temperature and they are based on a truncated log-logistic distribution of germination time:

$$P(t) = \frac{d}{1 + \exp[b(\log(x) - \log(e))]}$$

where the usual parameters ('b', 'd' and 'e') are expressed as functions of wtemperature (T). In the function 'TTEM()', we implemented the following submodels: (1) for the parameter 'd', we implemented the following submodels:

$$d = G \left[1 - \exp\left(-\frac{T_c - T}{\sigma_{T_c}}\right) \right]$$

$$1/[e(T)] = GR_{50}(T) = \frac{T - T_b}{\theta_T} \left[1 - \frac{T - T_b}{T_c - T_b} \right]$$

while 'b' was regarded as constant and independent from temperature.

In the 'TTERF()' function, the last submodel was modified, according to Rowse and Finch-Savage (2003):

$$1/[e(T)] = GR_{50}(T) = \begin{cases} \frac{T-T_b}{\theta_T} & \text{if } T_b < T < T_d \\ \frac{T-T_b}{\theta_T} \left[1 - \frac{T-T_d}{T_c-T_d} \right] & \text{if } T_d < T < T_c \\ 0 & \text{if } T \leq T_b \text{ or } T \geq T_c \end{cases}$$

In the 'TTERFc()' function a further submodel was introduced, to model the effect of temperature in the shape parameter 'b':

$$\sigma(T) = \frac{1}{b} = \frac{1}{b_0} + s(T - T_b)$$

The 'TTEM.fun()', 'TTERF.fun()' and 'TTERFc.fun()' are generic functions, which can be used for plotting or other applications, while the 'TTEM()', 'TTERF()' and 'TTERFc()' functions are meant to be used for model fitting with the 'drmt()' function in the 'drcte()' package.

Usage

```
TTEM()
TTEM.fun(time, Temp, G, Tc, sigmaTc, Tb, ThetaT, b)
TTERF()
TTERF.fun(time, Temp, G, Tc, sigmaTc, Td, Tb, ThetaT, b)
TTERFc()
TTERFc.fun(time, Temp, G, Tc, sigmaTc, Td, Tb, ThetaT, b0, s)
```

Arguments

The TTEM(), TTERF() and TTERFc() functions have no arguments. The TTEM.fun(), TTERF.fun() and TTERFc.fun() functions have the following arguments:

time	time
Temp	temperature
G	maximum germination capability of the seed lot
Tc	ceiling temperature
sigmaTc	standard deviation for the ceiling temperature within the seed lot
Tb	base temperature
Td	close to optimal temperature
ThetaT	Thermal-time parameter
b	shape parameter for the cumulative distribution function of germination time
b0	shape parameter for the cumulative distribution function of germination time at T = Tb
s	parameter for the effect of temperature on the scale parameter for the log-logistic distribution of germination times

Details

The detail of these functions are described in Onofri et al. (2018).

Value

The 'TTEM.fun()', 'TTERF.fun()' and 'TTERFc.fun()' functions return the proportion of germinated seeds, for given values of time and temperature. The 'TTEM()', 'TTERF()' and 'TTERFc()' functions return a list containing the nonlinear function, the self starter function, the parameter names and other items which are internally used by the 'drnte()' function.

Author(s)

Andrea Onofri

References

Onofri, A., Benincasa, P., Mesgaran, M.B., Ritz, C., 2018. Hydrothermal-time-to-event models for seed germination. *European Journal of Agronomy* 101, 129–139. Rowse, H.R., Finch-Savage, W.E., 2003. Hydrothermal threshold models can describe the germination response of carrot (*Daucus carota*) and onion (*Allium cepa*) seed populations across both sub- and supra-optimal temperatures. *New Phytologist* 158, 101–108. https://www.statforbiology.com/2023/stat_drcte_11-examplelette/

Examples

```
data(barley)
modTTE <- drnte(nSeeds ~ timeBef + timeAf + Temp, data = barley,
               fct = TTERF())
summary(modTTE)
```

Index

- * **capability of germination**
 - Pmax_models, 38
- * **datasets**
 - alfalfaSG, 2
 - barley, 4
 - excalibur, 5
 - festuca, 6
 - hordeum, 11
 - phalaris, 36
 - rape, 39
 - rape2G, 40
- * **germination rate**
 - GR_T_models, 10
 - GR_TPsi_models, 8
- * **hydro-thermal-time model**
 - HTTEM, 25
 - HTLL, 27
 - HTTnorm.M, 29
- * **hydro-time model**
 - HTE1, 13
 - HTE2, 14
 - HTE3, 16
 - HTex, 18
 - HTG, 19
 - HTL, 21
 - HTLL, 22
 - HTnorm, 24
 - HTW1, 31
 - HTW2, 32
- * **hydrotime models**
 - excalibur, 5
 - festuca, 6
 - hordeum, 11
 - rape, 39
 - rape2G, 40
- * **seed germination**
 - excalibur, 5
 - festuca, 6
 - GR_T_models, 10
 - GR_TPsi_models, 8
 - hordeum, 11
 - HTE1, 13
 - HTE2, 14
 - HTE3, 16
 - HTex, 18
 - HTG, 19
 - HTL, 21
 - HTLL, 22
 - HTnorm, 24
 - HTTEM, 25
 - HTTLL, 27
 - HTTnorm.M, 29
 - HTW1, 31
 - HTW2, 32
 - jackGroupSE, 34
 - makeDrm, 35
 - makeDrm2, 35
 - Pmax_models, 38
 - rape, 39
 - rape2G, 40
 - Thermal-Time_models, 41
- * **thermal-time model**
 - Thermal-Time_models, 41
- * **time-to-event model**
 - HTE1, 13
 - HTE2, 14
 - HTE3, 16
 - HTex, 18
 - HTG, 19
 - HTL, 21
 - HTLL, 22
 - HTnorm, 24
 - HTTEM, 25
 - HTTLL, 27
 - HTTnorm.M, 29
 - HTW1, 31
 - HTW2, 32
 - Thermal-Time_models, 41

alfalfaSG, 2

barley, 4

excalibur, 5

festuca, 6

GR_Psi_models, 7

GR_T_models, 10

GR_TPsi_models, 8

GRPsi.Lin (GR_Psi_models), 7

GRPsi.Pol (GR_Psi_models), 7

GRPsi.Pol2 (GR_Psi_models), 7

GRPsiLin (GR_Psi_models), 7

GRPsiPol (GR_Psi_models), 7

GRPsiPol2 (GR_Psi_models), 7

GRT.BS (GR_T_models), 10

GRT.BSb (GR_T_models), 10

GRT.Ex (GR_T_models), 10

GRT.Exb (GR_T_models), 10

GRT.GH (GR_T_models), 10

GRT.GH2 (GR_T_models), 10

GRT.M (GR_T_models), 10

GRT.Mb (GR_T_models), 10

GRT.RF (GR_T_models), 10

GRT.RFb (GR_T_models), 10

GRT.YL (GR_T_models), 10

GRTPsi.BS (GR_TPsi_models), 8

GRTPsi.M (GR_TPsi_models), 8

hordeum, 11

HTE1, 13

HTE2, 14

HTE3, 16

HTex, 18

HTG, 19

HTL, 21

HTLL, 22

HTnorm, 24

HTTEM, 25

HTTLL, 27

HTTnorm.BS (HTTnorm.M), 29

HTTnorm.M, 29

HTW1, 31

HTW2, 32

jackGroupSE, 34

makeDrm, 35

makeDrm2, 35

phalaris, 36

Pmax_models, 38

PmaxPsi1 (Pmax_models), 38

PmaxT1 (Pmax_models), 38

rape, 39

rape2G, 40

Thermal-Time_models, 41

TTEM (Thermal-Time_models), 41

TTERF (Thermal-Time_models), 41

TTERFc (Thermal-Time_models), 41