

# Package ‘geovol’

May 8, 2026

**Type** Package

**Title** Geopolitical Volatility (GEOVOL) Modelling

**Version** 1.1

**Date** 2025-05-06

**Description** Simulation, estimation and testing for geopolitical volatility (GEOVOL) based on the global common volatility model of Engle and Campos-Martins (2023) <[doi:10.1016/j.jfineco.2022.09.009](https://doi.org/10.1016/j.jfineco.2022.09.009)>. GEOVOL is modelled as a latent multiplicative volatility factor with heterogeneous factor loadings. Estimation is carried out as a maximization-maximization procedure, where GEOVOL and the GEOVOL loadings are estimated iteratively until convergence.

**License** GPL (>= 2)

**Depends** R (>= 3.5.0), zoo

**URL** <https://sites.google.com/site/susanacamposmartins/>

**NeedsCompilation** no

**Repository** CRAN

**Date/Publication** 2025-05-07 12:30:09 UTC

**Author** Susana Campos-Martins [aut, cre]

**Maintainer** Susana Campos-Martins <[sccmartins@ucp.pt](mailto:sccmartins@ucp.pt)>

## Contents

geovol-package . . . . .	2
coef.geovol . . . . .	3
geovol . . . . .	4
geovolObj . . . . .	6
geovolSim . . . . .	7
geovolTest . . . . .	8

<b>Index</b>	<b>9</b>
--------------	----------

---

geovol-package

*Geopolitical Volatility (GEOVOL) Modelling*

---

## Description

Simulation, estimation and testing for geopolitical volatility (GEOVOL) based on the global common volatility model of Engle and Campos-Martins (2023) <doi:10.1016/j.jfineco.2022.09.009>. GEOVOL is modelled as a latent multiplicative volatility factor with heterogeneous factor loadings. Estimation is carried out as a maximization-maximization procedure, where GEOVOL and the GEOVOL loadings are estimated iteratively until convergence.

## Details

Package: GEOVOL  
Type: Package  
Version: 1.1  
Date: 2025-05-06  
License: GPL>=2

## Author(s)

Susana Campos-Martins, <https://sites.google.com/site/susanacamposmartins/>

Maintainer: Susana Campos-Martins

## References

Engle, R.F. and Campos-Martins, S. (2023) Measuring and hedging geopolitical risk, *Journal of Financial Economics* 147: 221–242. Available at [doi:10.1016/j.jfineco.2022.09.009](https://doi.org/10.1016/j.jfineco.2022.09.009).

## See Also

[geovol](#), [geovolSim](#), [geovolTest](#)

## Examples

```
set.seed(123)

## Simulate from a GEOVOL model (default):
eSim <- geovolSim(n = 1500, m = 30)

## Test for GEOVOL:
geovolTest(e = eSim^2-1)
```

```
## Estimate a GEOVOL model:
eEst <- geovol(e = eSim)
eEst
```

---

coef.geovol                      *Extraction functions for a 'geovol' object*

---

## Description

Extraction functions for objects of class 'geovol'.

## Usage

```
## S3 method for class 'geovol'
coef(object, as.zoo = TRUE, ...)
## S3 method for class 'geovol'
fitted(object, as.zoo = TRUE, ...)
## S3 method for class 'geovol'
logLik(object, ...)
## S3 method for class 'geovol'
nobs(object, ...)
## S3 method for class 'geovol'
print(x, n.extreme = 20, ...)
## S3 method for class 'geovol'
residuals(object, as.zoo = TRUE, ...)
## S3 method for class 'geovol'
summary(object, ...)
```

## Arguments

object	an object of class 'geovol'.
x	an object of class 'geovol'.
as.zoo	logical. If TRUE, then the returned result is of class <code>zoo</code> .
n.extreme	integer indicating the n.extreme most extreme GEOVOL events to be printed.
...	additional arguments.

## Value

coef:	numeric vector containing parameter estimates.
fitted:	fitted (squared) GEOVOL time series.
logLik:	log-likelihood (normal density).
nobs:	the number of observations used in the estimation.
print:	print of the estimation results.
residuals:	GEOVOL standardised residuals.
summary:	summary of estimation results.

**Author(s)**

Susana Campos-Martins

**References**

Engle, R.F. and Campos-Martins, S. (2023) Measuring and hedging geopolitical risk, *Journal of Financial Economics* 147: 221–242. Available at [doi:10.1016/j.jfineco.2022.09.009](https://doi.org/10.1016/j.jfineco.2022.09.009).

**See Also**

[geovol](#), [geovolSim](#), [zoo](#)

**Examples**

```
set.seed(123)

## Simulate from a GEOVOL model (default):
eSim <- geovolSim(n = 1000, m = 30)

## Test for GEOVOL
eTest <- geovolTest(e = eSim^2)

## Estimate a GEOVOL model:
geovolEst <- geovol(e = eSim)

## Print estimation results:
print(geovolEst)

## Extract coefficients:
coef(geovolEst)

## Extract and store GEOVOL:
sigma2Est <- fitted(geovolEst)

## Extract optimised log-likelihood values:
logLik(geovolEst)

## Extract and store GEOVOL standardised residuals:
etaEst <- residuals(geovolEst)
```

---

geovol

*Estimate GEOVOL and GEOVOL loadings*

---

**Description**

Maximum likelihood estimation of the GEOVOL and GEOVOL loadings. Estimation is carried out as a maximization-maximization procedure, where the latent multiplicative volatility factor and the factor loadings are estimated iteratively until convergence. Either a balanced or an unbalanced panel can be used. For each point in time  $t$ , GEOVOL is estimated using all the available values in the cross-section. The GEOVOL loadings are estimated for each time series.

**Usage**

```
geovol(e, initial.values = list())
```

**Arguments**

- e** matrix, multivariate time series or [zoo](#) object of volatility standardized residuals with zero mean and unit variance.
- initial.values** a list containing the initial parameter values passed on to the optimization routine ([optim](#) for the GEOVOL factor and loadings. If `list()`, the default, then the values are chosen automatically; **x** - NULL or numeric vector containing the  $n$  initial values for the GEOVOL factor,  $x_{\{t\}}$ ,  $t=1, \dots, n$ . **s** - NULL or numeric vector containing the  $m$  initial values for the GEOVOL loadings,  $s_{\{i\}}$ ,  $i=1, \dots, m$ .

**Value**

An object of class 'geovol'.

**Author(s)**

Susana Campos-Martins

**References**

Engle, R.F. and Campos-Martins, S. (2023) Measuring and hedging geopolitical risk, *Journal of Financial Economics* 147: 221–242. Available at [doi:10.1016/j.jfineco.2022.09.009](https://doi.org/10.1016/j.jfineco.2022.09.009).

**See Also**

[geovol](#), [geovolSim](#), [geovolTest](#), [optim](#),

**Examples**

```
set.seed(123)

## Simulate from a GEOVOL model (default):
eSim <- geovolSim(n = 1000, m = 30)

## Test for GEOVOL
geovolTest(e = eSim^2-1)

## Estimate a GEOVOL model:
geovolEst <- geovol(e = eSim)

## Print estimation results:
print(geovolEst)

## Extract coefficients:
coef(geovolEst)
```

```
## Extract and store GEOVOL:
sigma2Est <- fitted(geovolEst)

## Extract optimised log-likelihood values:
logLik(geovolEst)

## Extract and store GEOVOL standardised residuals:
etaEst <- residuals(geovolEst)
```

---

geovolObj

*Auxiliary function*


---

### Description

Auxiliary function used in the estimation of the GEOVOL model. Not intended for the average user.

### Usage

```
geovolObj(par, par.fixed, y, x, flag)
```

### Arguments

par	scalar with the value of the parameter being estimated. If <code>x=TRUE</code> , par contains the value of the GEOVOL factor at time t. Otherwise, it contains the value of the GEOVOL loading for time series i.
par.fixed	numeric vector with the values of the parameters being fixed during estimation. If <code>x=TRUE</code> , par.fixed contains the values of the estimated GEOVOL loadings from previous iteration. Otherwise, it contains the values of the estimated GEOVOL factor from previous iteration.
y	numeric vector, time series or <a href="#">zoo</a> containing the cross-section at time t if <code>x=TRUE</code> or the time series i if otherwise.)
x	logical. If <code>TRUE</code> , it estimates the value of the GEOVOL factor at time t. Otherwise, it estimates the GEOVOL loading for time series i.
flag	integer. If 0, returns a numeric vector with the values of the objective function; if 1 returns the the value of the objective function; if 2, it returns the GEOVOL scaled by the loading of series i, if <code>x=FALSE</code> .

### Value

The values of the objective function or GEOVOL.

### Author(s)

Susana Campos-Martins

### See Also

[geovol](#), [fitted.geovol](#), [residuals.geovol](#)

---

geovolSim                      *Simulate from a GEOVOL model*

---

### Description

Simulate from a GEOVOL model.

### Usage

```
geovolSim(n, m, innovations = NULL, x.sd = 0.5, as.zoo = TRUE, verbose = FALSE)
```

### Arguments

n	integer indicating the number of observations.
m	integer indicating the number of time series.
innovations	NULL or numeric vector with the innovations. If NULL, then standard normal innovations are generated with <a href="#">rnorm</a> .
x.sd	numeric scalar indicating the standard deviation of the normal random variable, v, used to simulate the squared GEOVOL factor, x, according to the formula $x = \exp(v)$ .
as.zoo	logical. If TRUE, then the returned result is of class <a href="#">zoo</a> .
verbose	logical, if TRUE, the simulated time series, e, the innovations and the squared GEOVOL factor, x, are also returned.

### Value

A matrix with the simulated values.

### Author(s)

Susana Campos-Martins

### See Also

[geovol](#), [geovolTest](#),

### Examples

```
set.seed(123)

## Simulate from a GEOVOL model with s.d. 0.5 (default):
eSim <- geovolSim(n = 1000, m = 30)

## Simulate from a GEOVOL model with s.d. 1:
eSim <- geovolSim(n = 1000, m = 30, x.sd = 1)
```

---

`geovolTest`*Test for GEOVOL*

---

**Description**

Compute the GEOVOL test statistic for examining the null hypothesis that the average correlation of the squared volatility standardized residuals (which should be uncorrelated) is zero against the alternative that it is positive.

**Usage**

```
geovolTest(e)
```

**Arguments**

`e` matrix, multivariate time series or `zoo` object.

**Value**

The average correlation, the test statistic and the p-value.

**Author(s)**

Susana Campos-Martins

**References**

Engle, R.F. and Campos-Martins, S. (2023) Measuring and hedging geopolitical risk, Journal of Financial Economics 147: 221–242. Available at [doi:10.1016/j.jfineco.2022.09.009](https://doi.org/10.1016/j.jfineco.2022.09.009).

**See Also**

[geovol](#), [\]geovolSim](#)

**Examples**

```
set.seed(123)

## Simulate from a GEOVOL model with s.d. 0.5 (default):
eSim <- geovolSim(n = 1500, m = 30)

## Test for GEOVOL:
geovolTest(e = eSim^2-1)
```

# Index

## \* Econometrics

- coef.geovol, 3
- geovol, 4
- geovol-package, 2
- geovolObj, 6
- geovolSim, 7
- geovolTest, 8

## \* Financial Econometrics

- coef.geovol, 3
- geovol, 4
- geovol-package, 2
- geovolObj, 6
- geovolSim, 7
- geovolTest, 8

## \* Multivariate Time Series

- coef.geovol, 3
- geovol, 4
- geovol-package, 2
- geovolObj, 6
- geovolSim, 7
- geovolTest, 8

## \* Volatility Factor Models

- coef.geovol, 3
- geovol, 4
- geovol-package, 2
- geovolObj, 6
- geovolSim, 7
- geovolTest, 8

coef.geovol, 3

fitted.geovol, 6

fitted.geovol (coef.geovol), 3

geovol, 2, 4, 4, 5–8

geovol-package, 2

geovolObj, 6

geovolSim, 2, 4, 5, 7, 8

geovolTest, 2, 5, 7, 8

logLik.geovol (coef.geovol), 3

nobs.geovol (coef.geovol), 3

optim, 5

print.geovol (coef.geovol), 3

residuals.geovol, 6

residuals.geovol (coef.geovol), 3

rnorm, 7

summary.geovol (coef.geovol), 3

zoo, 3–8